arXiv:math-ph/0106027v2 16 Dec 2001

Poincaré normal and renormalized forms

Giuseppe Gaeta*
Dipartimento di Matematica, Universitá di Milano
V. Saldini 50, I–20133 Milano (Italy)
gaeta@roma1.infn.it

13 December 2001

Summary. We briefly review the main aspects of (Poincaré-Dulac) normal forms; we have a look at the non-uniqueness problem, and discuss one of the proposed ways to "further reduce" the normal forms. We also mention some convergence results.

1 Poincaré normal forms and their use

Normal forms were introduced by Poincaré as a tool to integrate nonlinear systems; by now we know this is in general impossible, but it turned out that the usefulness of normal forms goes well beyond integrability.

An introduction to normal forms is provided e.g. in [3, 5, 26, 41, 43, 47, 63]; see also [10, 33]. An introduction to normal forms for Hamiltonian systems is given in appendix 7 of [4].

Their use is, beyond integrability issues, of three kinds:

- Classification of dynamical systems;
- Qualitative study of dynamical systems;
- Quantitative study of dynamical systems.

In this note I want to discuss the Poincaré-Dulac approach, and an extension thereof, focusing on the simpler setting, i.e. normal forms for a smooth dynamical system (equivalently, an autonomous first order ODE, a vector field) in $\mathbf{R}^n$, around an equilibrium point. For more general settings, see e.g. [3, 5].

I consider a first order ODE (dynamical system) in R^n having a fixed point in the origin, and written as a series of homogeneous polynomial terms:

$$\dot{x} \;=\; Ax \,+\, \sum_{k=1}^{\infty} f_k(x) \tag{1}$$

*Paper to appear in *Acta Applicandae Mathematicae*; revised version. Work supported in part by "Fondazione CARIPLO per la ricerca scientifica" under project "Teoria delle perturbazioni per sistemi con simmetria".

where $f_k(ax) = a^{k+1} f_k(x)$ (we say also $f_k \in V_k$, and write $V = \oplus_{k=0}^{\infty} V_k$). We can of course think this is obtained from a general smooth[1] equation $\dot{x} = f(x)$, with $f(0) = 0$, expanding it[2] around $x = 0$. Actually, for most of our discussion, we could as well allow the series on the r.h.s. of (1) to be just a formal series (i.e. not require convergence in any neighbourhood of the origin).

I will also *assume that A is semisimple*[3]; for a discussion of the general case, see [47, 65] and [5, 19, 33]. Recall that if A is semisimple, it can be brought to diagonal form, and thus to be normal, $[A, A^+] = 0$ (this will be the relevant property); it will thus be more economic to perform all computations in a basis where A is diagonal.

The idea of Poincaré was to eliminate nonlinear terms by a sequence of near-identity changes of coordinates,

$$x \;=\; y \,+\, h_m(y) \tag{2}$$

with $h_m \in V_m$, and m taking successively the values $m = 1, 2, ...$

With this, and writing D for the matrix of partial derivatives of h_m, eq.(1) reads $[I + D]\dot{y} = f(y + h_m(y))$, which means

$$\dot{y} \;=\; [I + D]^{-1} \sum_{k=0}^{\infty} f_k(y + h_m(y)) \;:= \sum_{k=0}^{\infty} \widetilde{f}_k(y) \tag{3}$$

By a simple computation, we obtain that $\widetilde{f}_k(x) = f_k(x)$ for $k < m$, and (see below for the notation)

$$\widetilde{f}_m(x) \;=\; f_m(x) - \mathcal{L}_0(h_m) \;; \tag{4}$$

the effect on higher order terms is more involved, and is usually not considered in detail, as higher order terms can be normalized at later stages.

In eq.(4) above, $\mathcal{L}_0$ is a linear operator, defined in terms of the linear part A of f as

$$\mathcal{L}_0(\psi) \;:=\; (Ax \cdot \nabla)\psi \,-\, A\psi \;; \tag{5}$$

notice that $\mathcal{L}_0 : V_k \to V_k$. Actually it is more convenient to describe this is terms of a general bracket

$$\{\varphi, \psi\} \;:=\; (\varphi \cdot \nabla)\psi \,-\, (\psi \cdot \nabla)\varphi \tag{6}$$

(this is just the representation of the commutator of the vector fields $\varphi^i \partial_i$ and $\psi^i \partial_i$ in terms of their components); with this, $\mathcal{L}_0$ is just defined as $\mathcal{L}_0(\psi) := \{Ax, \psi\}$.

[1] By this we will always mean C^∞.

[2] In this way we lose all the information "beyond all orders" in perturbation theory, such as e^{-1/x^2} terms; for this, see the works by Ecalle. Notice that if we deal with perturbative power series expansions of smooth dynamical systems (vector fields), it would be natural in some aspect to consider C^∞ rather than analytic conjugacy.

[3] Let me stress, however, that a good part of the discussion conducted here will also apply to the case where A is not semisimple (at least provided it has nonzero semisimple part).

In this way we can eliminate all the terms which are in the range of $\mathcal{L}_0$ (note that if A is normal, $\mathrm{Ker}(\mathcal{L}_0)$ is complementary to $\mathrm{Ran}(\mathcal{L}_0)$ in V, and thus the whole normal form can be taken to be in $\mathrm{Ker}(\mathcal{L}_0)$; this means that the whole normal form commutes with the linear part of the system). We can also introduce a scalar product[4] in V, so that $\mathrm{Ran}(\mathcal{L}_0) = \mathrm{Ker}(\mathcal{L}_0^+)$.

We can now also easily characterize (with A taken to diagonal form) the terms in $\mathrm{Ker}(\mathcal{L}_0^+)$; assume $\lambda_1, ..., \lambda_n$ are the eigenvalues of A, and $\mathbf{v}_1, ..., \mathbf{v}_n$ the corresponding eigenvectors (the computations are immediate if we take a basis of eigenvectors in R^n). Then a vector $\mathbf{w} = x^\mu \mathbf{v}_r := x_1^{\mu_1} ... x_n^{\mu_n} \mathbf{v}_r$ is in $\mathrm{Ker}(\mathcal{L}_0^+)$ if and only if there are nonnegative integers $\mu_1 ... \mu_n$ (with $|\mu| := \sum_{s=1}^n \mu_s > 1$) such that

$$\sum_{s=1}^{n} \mu_s \lambda_s \;=\; \lambda_r \tag{7}$$

and $\mathrm{Ker}(\mathcal{L}_0^+)$ is spanned by such "resonant monomials". The equation (7) is called a *resonance relation* of order $|\mu|$.

Thus, following the Poincaré procedure, we can always eliminate term by term all the nonlinear terms up to any desired order N, provided the λ_i are nonresonant; note that this is generically the case.

In the case where the λ_i satisfy some resonance relation, i.e. when (7) admits some solution, we can always eliminate nonresonant terms, and arrive at a form of f, the *Poincaré-Dulac normal form*, in which only resonant terms are present. That is, we arrive at coordinates in which (1) is written as[5]

$$\dot{x} \;=\; Ax + \sum_{k=1}^{\infty} g_k(x) \;\;;\;\; g_k \in \mathrm{Ker}(\mathcal{L}_0^+) \; . \tag{8}$$

It should be stressed that this result also holds for non-semisimple A [33].

In this way we obtain a description of nonlinear systems (in normal form) "compatible" with a given linear part.

In practice, the normalization will be carried out only up to some finite order N; one will then use perturbative arguments to infer that the dynamics of the full system is only a small modification – at least in a sufficiently small neighbourhood of the origin (and/or for sufficiently short times) – of that described by the truncated normal form. In this way, if we were able to study the most general (truncated) system in the form (8) for given A, we would in principles obtain a classification of the qualitative local behaviours of nonlinear systems around a fixed point with prescribed linearization A.

Let me point out, after mentioning these "qualitative" uses of normal forms, the basis for its use in quantitative studies. Consider again (1), and denote by μ the first k with $f_k \not\equiv 0$ (so generically $\mu = 1$); rescale x as $x = \varepsilon \widetilde{x}$, divide by

[4] With the notation introduced below, the scalar product in V can be chosen to be the Bargmann scalar product [47], $(x^\mu \mathbf{v}_r, x^\nu \mathbf{v}_s) := \delta_{r,s} \delta_{\mu,\nu} (\mu!)$. With this, $\mathcal{L}_0^+(\psi) = \{A^+ x, \psi\}$.

[5] More precisely, we can arrive to such a form up to order N, for any desired N. That is, we should add a remainder $\mathcal{R}_N$ of arbitrarily high order $N+1$ in this equation.

ε, and drop the tilde for ease of notation, so that we have a system of the form

$$\dot{x} \;=\; Ax \,+\, |\varepsilon|^{\mu} g(x;\varepsilon)\;; \tag{9}$$

let $x(t) = \Phi_\varepsilon(x_0, t)$ be its solution with initial datum $x(0) = x_0$. If we approximate this with the solution $\xi(t) = \Phi_0(x_0, t)$ to the system $\dot{x} = Ax$ obtained for $\varepsilon = 0$, we are making an error

$$\rho_\varepsilon(t) \;:=\; |x(t) - \xi(t)| \;=\; |\Phi_\varepsilon(x_0, t) \,-\, \Phi_0(x_0, t)| \tag{10}$$

Assume that A has norm C, and that in a region $\mathcal{B}$ we have $|g(x;\varepsilon)| \le M(\varepsilon)$ (all of our estimates will be only valid when x and ξ are both in $\mathcal{B}$, i.e. for a time t depending on the exit time from $\mathcal{B}$). Then one obtains easily[6] that

$$\rho_\varepsilon(t) \;\le\; |\varepsilon|^{\mu} \frac{M}{C} \left[e^{Ct} - 1\right] \;. \tag{11}$$

This also means that $\rho_\varepsilon(t) < \delta$ for a time not smaller than

$$t_0 \;=\; \frac{1}{C} \;\ln\left[1 + \frac{\delta C}{|\varepsilon|^{\mu} M}\right] \;. \tag{12}$$

When we perform a normalization, with no resonance of order smaller than μ, we are pushing nonlinearities to higher orders and thus increasing μ in the previous estimates; that is, we are improving the accuracy of the linear estimate, or extending the time interval over which we have an error of the same "acceptable" size δ.

It should be stressed that in changing coordinates we are also changing the value of the constant M; if the "small denominators" appearing due to the change of coordinates are too small, their effect will more than counterbalance the gain we obtain by pushing up the nonlinearities, and disrupt the perturbative expansion.

2 The problem of convergence

It should be noted that the discussion conducted above is purely formal: the changes of coordinates are constructed as a series which could – and in general will – be only formal.

Here we want to consider the case of analytic systems, and for these we should study the convergence of the series defining the transformations.

This is in general a hard problem (let us stress that it is difficult to prove convergence, but it is also difficult to prove divergence; see [3, 5]), and there are very few general results. Let me recall the main ones[7] [3, 26, 27, 65]. Here

[6] Indeed $d\rho_\varepsilon/dt \le |A(x-\xi)| + |\varepsilon^{\mu} g(x,\varepsilon)| \le C\rho_\varepsilon + |\varepsilon|^{\mu} M(\varepsilon)$. Thus the growth of $\rho_\varepsilon(t) > 0$ is slower than that of a function $r(t)$ obeying $dr/dt = ar + b$ with $r(0) = 0$ and $a = C$, $b = |\varepsilon|^{\mu} M(\varepsilon)$. The solution to this equation is $r(t) = (b/a)[\exp(at) - 1]$.

[7] It is remarkable that when we can guarantee convergence of the normalizing transformation on the basis of the first four criteria below, we have an *integrable normal form.*

the function $f : \mathbf{R}^n \to \mathbf{R}^n$ defining the dynamical system $\dot{x} = f(x)$ should be thought of as $f : \mathbf{C}^n \to \mathbf{C}^n$.

• *Poincaré criterion:* if the convex hull of the $\lambda_i \in \mathbf{C}$ does not include zero, then the normalizing series converges in an open neighbourhood of the origin.

• *Siegel criterion:* call the vector $\lambda = (\lambda_1, ..., \lambda_n)$ of type (C, ν) if for each $r = 1, ..., n$ and for any $|\mu| \geq 2$ it satisfies $|\sum_s (\mu_s \lambda_s) - \lambda_r| \geq C/|\mu|^\nu$. If f is analytic and λ is of type (C, ν) for some positive numbers C and ν, then f is analytically equivalent to its linear part. (Note that the set of vectors λ which are not of type (C, ν) for any $C > 0$ has measure zero for $\nu > (n-2)/2$; in the real case this holds for $\nu > n - 1$).

• *Pliss criterion:* Pliss realized that Siegel condition can be weakened: call the vector $\lambda = (\lambda_1, ..., \lambda_n)$ of type $P(K, \nu)$ if for each $r = 1, ..., n$ and for any $|\mu| \geq K$, it satisfies either the resonance condition $|\sum_s (\mu_s \lambda_s) - \lambda_r| = 0$, or $|\sum_s (\mu_s \lambda_s) - \lambda_r| \geq 1/|\mu|^\nu$. If f is analytic, λ is of type $P(K, \nu)$ for some positive numbers K and ν, and f is formally conjugated to its linear part, then f is analytically conjugated to its linear part.

• *Bruno criterion:* say that λ satisfies "condition ω" if $\sum_{k=1}^{\infty} 2^{-k} \ln(1/\omega_k) < \infty$, where ω_k is the minimum of $|\sum_s \mu_s \lambda_s - \lambda_r|$ on all the $r = 1, ..., n$ and all the $|\mu| < 2^k$ not satisfying a resonance relation (condition ω is weaker than the Siegel or Pliss condition [19]). Say that f having linear part Ax satisfies "condition A" if[8] $f(x) = [1 + \alpha(x)]Ax$. Then, if λ satisfies condition ω and f is formally conjugated to a normal form $\widehat{f}$ satisfying condition A, then f is conjugated to $\widehat{f}$ by an analytic transformation. (Condition A can be weakened for suitable classes of f's).

• *BMW-C theory:* assuming condition ω for λ, condition A can be replaced by suitable symmetry conditions on f. This theory was created by Markhashov [52] and by Bruno and Walcher [21]; it was later extended by Cicogna [25] and again by Walcher [69]. This matter is discussed in [27] and in the SPT2001 tutorial by Cicogna and Walcher [28]. See also the recent work by Stolovich [59] which, together with many other things, contains results in this direction.

• *Normalization of an algebra of vector fields:* one can consider the problem of simultaneously normalizing an algebra of (analytic) vector fields X_i around a common singular point. Vey [64] proved that if they span an abelian algebra and their linearizations at the singular point are linearly independent, then they are simultaneously and analytically normalizable. Results in this direction are also contained in [2, 42, 60]; see also [26].

• *Hamiltonian systems:* in the case of Hamiltonian systems, we can deal directly with the Hamiltonian (i.e. a scalar function) rather than with the hamiltonian vector field this generates. Many of the results quoted above can be recast in this setting, but there have been investigations of the convergence in the hamiltonian case *per se*. In this context, we refer to the work of Russmann, Bruno, Vey, and Ito [19, 48, 56, 64], as well as to that of Giorgilli and collaborators [38] (also in connection with the study of effective stability and

[8] In geometrical terms, if f is orbitally equivalent to its linear part [67]. This means the trajectories of f are the same as those of Ax.

KAM theory [39, 40]). I will not discuss these results here; see however the paper by Cicogna in [9] and [26, 27] for brief discussions.

• *BCS theory:* convergence of the normalizing transformation implies that the original vector field and its normal form are analytically conjugated. In many cases one would be satisfied with C^∞ conjugation [10, 13] (or even with C^k conjugation: e.g., this is the only one making sense for C^k vector fields. For C^k normal forms theory, see [46]). It was proved by Chen and by Sternberg [22, 58] that if the linear part A of the system is hyperbolic, then formal conjugacy implies C^∞ conjugacy (see also [10, 30, 44]). This result has been recently extended to equivariant (or reversible) systems, and also to the Hamiltonian framework, by Belitskii and Kopanskii [14].

• *Orbital conjugacy:* the reader should be warned that in many cases one is satisfied with orbital equivalence (two systems are orbitally equivalent if they have the same integral curves, independently of the time law they follow on these). This question is considered e.g. in [5, 45, 67].

In the following, I will come back to considering only formal transformations; for a discussion of convergence see e.g. [26, 27, 28].

3 Some remarks on Poincaré normalization

Now that I described the why and how of Poincaré normalization, let me remark some simple points which are relevant for the following.

- The change of coordinates needed to normalize f at order k is generated by h_k solving $\mathcal{L}_0(h_k) = \pi_0(f_k)$, where π_0 is the projection from V to the range of $\mathcal{L}_0$, and f_k is the term obtained after the first $(k-1)$ normalizations. The solution to this "homological equation" for h_k is unique up to a term in $\mathrm{Ker}(\mathcal{L}_0)$. That is, if $\delta h_k = h_k - \widehat{h}_k \in \mathrm{Ker}(\mathcal{L}_0)$, then h_k and $\widehat{h}_k$ define the same transformation in V_k.

- On the other hand, functions h_k and $\widehat{h}_k$ as above have in general a different effect on the f_p with $p > k$. Thus the freedom in the choice of h_k (for resonant systems) means that we can generate different normal forms starting from the same nonlinear system.

- This shows at once that normal forms are somehow ambiguous, i.e. the normal forms classification is in general redundant: different normal forms (with the same linear part) can be actually conjugated.

- It is maybe worth remarking explicitely that the higher order terms generated by the change of coordinates $x = y + h_k(y)$ will in general also have a resonant part; this cannot be eliminated by the Poincaré procedure at later stages[9].

[9] Thus, in particular for resonant systems, it is possibly misleading to say that normalization is a way to "simplify" (a rather vague word) nonlinear systems: a system with very few terms can have infinitely many terms once taken into normal form.

It is clear that from the point of view of qualitative study and classification we would like to eliminate or at least reduce the redundancy in the normal forms classification. This would also have an obvious computational advantage: among normal forms corresponding to f we could choose the most convenient for our needs.

Thus the problem of "further normalization" has been considered by several authors[10]; I will quote first of all Takens [61], and Broer who in his thesis (advised by Takens) set the problem – and the theory of normal forms – in terms of filtrations of Lie algebras [16, 17, 18]. In this respect, one should also mention the early work of Kummer [51].

Broer's Lie algebra approach was also followed by van der Meer [53] and by Baider, Churchill and Sanders [6, 7, 8]; in particular they obtained a formulation of "unique normal forms", see also [24, 50]. Unfortunately, their results are very difficult to implement on concrete systems, even the simpler ones, and seems to be not applicable in concrete terms (on the other hand, Broer's approach is a guiding light for many questions in normal forms theory).

Other, and more recent, attempts in this direction (some of them with different tools) include works by Algaba, Freire and Gamero, by Chen and Della Dora, by Kokubu, Oka and Wang, by Ushiki, and by myself [1, 23, 50, 35, 62].

Unfortunately, no systematic exposition and comparison of these different approaches is available; needless to say, this is not the place to attempt such a work, but maybe some reader of the present paper will endeavour a comprehensive discussion of this matter, which would surely be useful.

The same problem of further normalization was also tackled, of course, by russian authors. One should quote in this respect at least the works by Belitskii [12] and by Bruno [19]; these are also discussed and compared in the recent book [20]. For russian work in the area see also, of course, [5].

In the next sections I will present a simple way[11] to (in general, only partially) remove the degeneration of normal forms. This is based on ideas which I believe were first introduced by Broer (and recalled in a somewhere deformed way in the final section 6); the approach discussed below in sections 4 and 5 aims at a smaller generality than Broer's one, but is easily implemented in practice [35, 36].

I would like to thank prof. Belitskii for pointing out, after a preliminary version of the present paper was circulated, that this approach is very much similar to the one proposed in his work [11].

[10] Actually already Dulac (see page 335 of [32]) remarked that possibly a choice of the terms δh_k in $\mathrm{Ker}(\mathcal{L}_0)$ different from the one he was employing – i.e. $\delta h_k = 0$ – could have produced a simpler normal form.

[11] The reader should be warned that the discussion of it given in [20] uses different basic definitions; see [36].

4 Renormalization (further normalization)

The approach I will discuss in this section (based on [35]; see also [26], and [36] for more recent discussions) is completely algorithmic, and not any more difficult to implement (by hand or by computer) than the standard Poincaré one: actually it is just based on a careful exploitation of the freedom left by Poincaré algorithm. As mentioned above, this is quite similar to the one proposed in [11].

The main idea is to control the effects of the change of variables generated by h_k on higher order terms f_p, $p > k$; as already mentioned, this change of variables will in general produce resonant terms. This should not be seen as a drawback, but rather as an advantage! By a judicous choice of h_k, we can use this feature to *eliminate higher order resonant terms*. This "judicious choice" amounts to choosing the h_k's as solutions to "higher homological equations".

It is actually convenient to consider an improved version of Poincaré transformations, i.e. Lie-Poincaré ones; for some of the advantages of using these see [15], see also [31, 66]. The idea is to define a vector field

$$H \;:=\; h^i_m(x)\,\frac{\partial}{\partial x^i} \tag{13}$$

in $\mathbf{R}^n$ and consider the flow $\Phi(\sigma; x_0)$ generated by this. The change of coordinates is then given by $x = \Phi(1; y)$, which coincides with the standard one at lowest nontrivial order.

The change of coordinates with generator h_m maps (1) into a new system $\dot{y} = \widehat{f}(y) = Ay + \sum_{k=1}^{\infty} \widehat{f}_k(y)$, where $\widehat{f}_k(x) = f_k(x)$ for $k < m$, while for $k \geq m$ we have, with $[a]$ the integer part of a,

$$\widehat{f}_k(x) \;=\; \sum_{s=0}^{[k/m]} \frac{1}{s!}\,\mathcal{H}^s_m(f_{k-sm})\;; \tag{14}$$

here the operator $\mathcal{H}_m$ is defined as $\mathcal{H}_m(\alpha) := \{h_m, \alpha\}$, and $\mathcal{H}^p_m$ means applying $\mathcal{H}_m$ for p times. Note that the expression (14) is just the translation to the present notation of the familiar Baker-Campbell-Hausdorff formula: indeed it states that the vector field $X = f^i\partial_i$ is changed into a vector field $\widehat{X} = \widehat{f}\widehat{\partial}_i$ given by $\widehat{X} := \left[e^{\sigma H} X e^{-\sigma H}\right]_{\sigma=1}$.

Having determined the effect of the Lie-Poincaré transformation to all orders, we can try to control it.

Let us consider again our system (1), i.e. $\dot{x} = f(x) = Ax + \sum_{k=1}^{\infty} f_k(x)$. We can normalize the quadratic term f_1 by solving the homological equation $\mathcal{L}_0(h_1) = \pi_0 f_1$, and we get[12]

$$\widehat{f}_1 \equiv f^{(1)}_1 = f_1 - \mathcal{L}_0(h_1)\;. \tag{15}$$

The higher order terms f_k ($k \geq 2$) will of course also change with this first change of variables, but for the sake of notation we will not introduce indices to stress this fact, and call the new expressions of these again $f_2, f_3, ...$

[12]The notation $\widehat{f}_1$ points out that this term is stable, i.e. the algorithm will not change it any more.

Now, consider the cubic term f_2; we normalize it by solving $\mathcal{L}_0(h_2) = \pi_0 f_2$, and we get $f_2^{(1)} = f_2 - \mathcal{L}_0(h_2)$. If this is nonzero we would like to (attempt to) further eliminate some terms. We can do it by exploiting the higher order effects of a transformation with generator $h_1^{(1)}$, and in order to avoid this affects $f_1^{(1)}$ we have to require $h_1^{(1)} \in \mathrm{Ker}(\mathcal{L}_0)$. From the general formula (14) we have then that with such a transformation

$$\widehat{f_2} \equiv f_2^{(2)} = f_2^{(1)} + \{h_1^{(1)}, f_1^{(1)}\} . \tag{16}$$

We define the linear "higher homological operators" $\mathcal{L}_k$ as

$$\mathcal{L}_k(\alpha) := \{\widehat{f_k}, \alpha\} \tag{17}$$

so that the above reads

$$f_2^{(2)} = f_2^{(1)} - \mathcal{L}_1\left(h_1^{(1)}\right) , \tag{18}$$

and we can eliminate from $f_2^{(1)}$ all terms which are in the range of $\mathcal{M}_1$, defined as the restriction of $\mathcal{L}_1$ to $\mathrm{Ker}(\mathcal{L}_0)$. Thus, if we choose the complementary spaces to the ranges considered as the orthogonal complement, we get $\widehat{f_2} \in [\mathrm{Ran}(\mathcal{L}_0)]^\perp \cap [\mathrm{Ran}(\mathcal{M}_1)]^\perp$.

To avoid any misunderstanding, we stress immediately that our procedure is based on these restricted operators $\mathcal{M}_k$, see below, and *not* on the full higher homological operators $\mathcal{L}_k$.

Higher order terms, f_3 and so on, will also change; again we will denote them, for the sake of notation, still as f_3 etc. not to introduce heavy notation.

Let us, for the sake of clarity, go one step further and consider the quartic term f_3. First we normalize it by a transformation generated by h_3, obtaining a $f_3^{(1)} \in \mathrm{Ker}(\mathcal{L}_0^+)$. Then we can still use a transformation generated by $h_2^{(1)} \in \mathrm{Ker}(\mathcal{L}_0)$ to eliminate some terms without affecting f_2; we obtain in this way a $f_3^{(2)}$. But we can also still use a transformation generated by $h_1^{(3)}$ provided this does not affect $\widehat{f_1}$ nor $\widehat{f_2}$, i.e. provided $h_1^{(3)} \in \mathrm{Ker}(\mathcal{L}_0) \cap \mathrm{Ker}(\mathcal{L}_1)$; with this, we obtain

$$\widehat{f_3} = f_3^{(2)} - \mathcal{L}_2\left(h_1^{(3)}\right) , \tag{19}$$

i.e. we can eliminate also all the terms which are in the range of $\mathcal{M}_2$, the restriction of $\mathcal{L}_2$ to $\mathrm{Ker}(\mathcal{L}_0) \cap \mathrm{Ker}(\mathcal{L}_1)$.

At this point the general procedure should be quite clear; in order to state precisely the result I should define some chains of spaces and operators[13].

• Let $H^{(0)} := V$, and define $H^{(p+1)} := \mathrm{Ker}(\mathcal{L}_0) \cap ... \cap \mathrm{Ker}(\mathcal{L}_p)$. Notice that $H^{(p+1)} \subseteq H^{(p)}$.

[13]Note that all of these spaces could be intersected with, and operators restricted to, the V_k, obtaining finite dimensional spaces and operators [35]. I will not do this to avoid a cumbersome notation, but in computer algebra implementation of this algorithm one would proceed in this way.

• Define the operators $\mathcal{M}_p := \mathcal{L}_p|_{H^{(p)}}$.
• Let $F^{(0)} := V$, and define $F^{(p+1)} := [\mathrm{Ran}(\mathcal{M}_0)]^\perp \cap ... \cap [\mathrm{Ran}(\mathcal{M}_p)]^\perp$. Notice that $F^{(p+1)} \subseteq F^{(p)}$.

It will also be convenient to set a notation for relevant projection operators:

• Denote by χ_s the projection operator $\chi_s : V \to \mathrm{Ker}(\mathcal{L}_s)$
• Denote by μ_s the composition of projection operators $\mu_s = \chi_{s-1} \cdot ... \cdot \chi_0$ for $s \geq 1$; we set $\mu_0 \equiv I$.
• Denote by π_s the projection operator $\pi_s : V \to \mathrm{Ran}(\mathcal{M}_s)$.

Note that with these definitions, $H^{(p)} = \mu_p V$, and $\mathcal{M}_p = \mathcal{L}_p \circ \mu_p$.

Definition. The dynamical system $\dot{x} = f(x) = \sum_{k=1}^{\infty} f_k(x)$ is in *Poincaré renormalized form* (PRF) up to order N if $f_k \in F^{(k)}$ for all $k < N$. In the formal limit $N \to \infty$, we say the system is in PRF, tout court.

Theorem. *Any analytic or formal smooth dynamical system can be taken into PRF up to any desired order N by means of a sequence of Lie-Poincaré transformations.*

Proof. Proceeding as above, the transformation generated by $h^{(p)}_{k-p}$ will take $f^{(p)}_k$ into

$$f^{(p+1)}_k \;=\; f^{(p)}_k \;-\; \mathcal{M}_p \left(h^{(p)}_{k-p} \right) \tag{20}$$

If we choose $h^{(p)}_{k-p}$ as a (non-unique) solution to the higher homological equation

$$\pi_p \, f^{(p)}_k \;-\; \mathcal{M}_p \left(h^{(p)}_{k-p} \right) \;=\; 0 \tag{21}$$

i.e. as

$$h^{(p)}_{k-p} \;=\; \mu_p \cdot \mathcal{M}^*_p \cdot \pi_p \left[f^{(p)}_k \right] \tag{22}$$

where $\mathcal{M}^*_p$ is the pseudo-inverse to $\mathcal{M}_p$, then we arrive – after applying the full procedure to $f_1, ..., f_N$ – to a system $\dot{x} = f^*(x) = \sum_{k=0}^{\infty} f^*_k(x)$ where $f^*_k(x) = f^{(k)}_k \in F^{(k)} \cap V_k$ for all $k \leq N$. △

Remark. The proof is completely constructive, as (22) tells which generator should be chosen at each step (and this is of course the nontrivial content of the theorem). It can thus be transformed in an algorithm to be implemented on a computer. ⊙
Remark. The algorithm can deal with systems having zero linear part, contrary to the standard normalization algorithm. For such an application in a symmetry context, see [37]. ⊙
Remark. The algorithm is only based on the grading of resonant terms (resonant vector fields) by their homogeneity properties, and is therefore completely general. However, it fails to recognize the Lie algebraic properties of the set of resonant vector fields. Not surprisingly, taking these into account leads to a more powerful reduction, which we discuss below. ⊙

Notice that this procedure, as discussed here, is completely formal. A discussion (certainly not final) of convergence of PRFs – and also of PRF for Hamiltonian systems, and of PRF in the presence of symmetry – is contained in [26] and in my contributions to the SPT workshops [9, 29]. A discussion of these matters is certainly incompatible with the limited scope of this paper.

I would now like to briefly mention an example of application of PRFs to a concrete problem, i.e. the (formal) analysis at singular points of vector fields in the plane having purely rotational linear part; this case corresponds to the vector field being, in the first order approximation, an integrable hamiltonian vector field. A complete analysis of regular singular points of vector fields in the plane from the point of view of PRFs is contained in [36], where analytic computations are described in detail, and completely explicit computations are pursued up to order six (order ten in a special case).

Example. Let the dynamical system $\dot{x} = f(x)$ in $\mathbf{R}^2$ have a singularity at the point x_0; we can always shift coordinates so that $x_0 = 0$. Let the linearization $\dot{x} = [(\partial_j f^i)(x_0)]x^j \equiv A^i_j x^j$ of X at $x_0 = 0$ be given by

$$A \;=\; \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \tag{23}$$

Then the standard normal form is given (in vector notation) by

$$\dot{x} \;=\; Ax \;+\; \sum_{k=1}^{\infty} r^{2k} \left(a_k I \;+\; b_k A\right) x \tag{24}$$

where I is the identity matrix and $r^2 = |x|^2 = (x_1^2 + x_2^2)$.

In general, a_1, b_1 are nonzero (but if the system is hamiltonian, all the a_k vanish; this case is treated in [34, 57]); however, for the sake of generality, let us assume $a_k = 0$ for $k < \mu$ and $b_k = 0$ for $k < \nu$, with $a_\mu \neq 0$ and $b_\nu \neq 0$.

Then, following step by step the procedure described above, the PRF is given by

$$\begin{cases} \dot{w} \;=\; Aw \;+\; (r^{2\mu}\alpha_\mu + r^{4\mu}\alpha_{2\mu})\, w & \text{for } \mu < \nu \\ \dot{w} \;=\; (1 + r^{2\mu}\beta_\mu)\, Aw \;+\; (r^{2\mu}\alpha_\mu + r^{4\mu}\alpha_{2\mu})\, w & \text{for } \mu = \nu \\ \dot{w} \;=\; (1 + r^{2\nu}\beta_\nu)\, Aw \;+\; \sum_{k=\mu}^{\infty}(r^{2k}\alpha_k)\, w & \text{for } \mu > \nu \end{cases} \tag{25}$$

where α_k, β_k are real numbers (in general different from the a_k, b_k).

This result was obtained in [35]; however the computation (and the result reported) there is not correct in the case with $\mu > \nu$; the correct computation leading to (25) is given in [36].

Notice that the standard NF is characterized by two infinite sequences of constants; the PRF is finite except in case $\mu > \nu$, where it however depends on only one infinite sequence of constants.

5 Lie renormalization

Let us look more closely at the set $\mathcal{G}$ of nonlinear vector fields resonant with a given linear one, $X_A = (Ax)^i \partial_i$.

First of all, it is clear that they form a Lie algebra (in general infinite dimensional) under commutation. Actually this also belongs to a finitely generated module, as we now briefly discuss recalling a general characterization of vector fields in normal form [26, 33, 47, 65].

We say that the differentiable function $\varphi : \mathbf{R}^n \to \mathbf{R}$ is an invariant for X_A if $X_A(\varphi) = 0$. Denote by $\mathcal{I}(A)$ the set of invariants for X_A which are meromorphic (that is, can be expressed as a quotient of analytic functions) in the x coordinates; and denote by $\mathcal{I}_*(A) \subset \mathcal{I}(A)$ the set of analytic invariants for X_A.

Let $G = C(A)$ be the centralizer of A in the algebra of n dimensional matrices; let its Lie algebra be spanned by matrices $\{K_1, ..., K_d\}$ (we can always assume $K_1 = I$, and take one of the K_α to be A if this is nonzero; notice that $d \leq n$). We denote by $X^{(\alpha)}$ the vector fields corresponding to these, i.e. given in the x coordinates by $X^{(\alpha)} = (K_\alpha x)^i \partial_i$.

Then it results [26, 33, 47, 65] that the most general vector field W in $\mathcal{G}$ can be written as

$$W \;=\; \sum_{\alpha=1}^{d} \mu_\alpha(x)\; X^{(\alpha)} \tag{26}$$

where $\mu_\alpha(x) \in \mathcal{I}(A)$; this means that $\mathcal{G}$ is contained in the module over $\mathcal{I}(A)$ generated by G.

Notice that the vector field W must be analytic in the x, while generic vector fields of the form $\mu_\alpha(x) X^{(\alpha)}$ are not such: $\mathcal{G}$ is not the full G-generated module over $\mathcal{I}(A)$.

In several cases it happens that $\mathcal{G}$ has a more convenient structure, i.e. the μ_α in (26) can actually be taken to be in $\mathcal{I}_*(A)$, and not just in $\mathcal{I}(A)$. In this case we have $\mathcal{G} = \mathcal{I}_*(A) \otimes G$, and we say that all the vector fields in $\mathcal{G}$ are *quasi-linear*, or that we have a **quasi-linear normal form**. In particular, this is e.g. the case when A admits only one basic invariant.

Call $\mathcal{X}_\alpha$ the algebra spanned by vectors which are written as $X = s(x) X^{(\alpha)}$ with $s \in \mathcal{I}(A)$; call $\mathcal{X}^*_\alpha \subset \mathcal{X}_\alpha$ the algebra spanned by vectors as above with $s \in \mathcal{I}_*(A)$ (this is the full module over $\mathcal{I}_*(A)$ generated by $X^{(\alpha)}$). As seen before, in general we have $\mathcal{X}^*_1 \oplus ... \oplus \mathcal{X}^*_d \subseteq \mathcal{G} \subset \mathcal{X}_1 \oplus ... \oplus \mathcal{X}_d$, and in the quasi-linear case we actually have $\mathcal{G} = \mathcal{X}^*_1 \oplus ... \oplus \mathcal{X}^*_d$.

Consider now the commutation relations between elements $\mu_\alpha(\Psi) X^{(\alpha)}$ and $\sigma_\beta(\Psi) X^{(\beta)}$ of the subalgebras $\mathcal{X}_\alpha$ and $\mathcal{X}_\beta$ respectively ; it is immediate to check that, with $\nabla_i := \partial / \partial \psi_i$, we have

$$\begin{aligned} \left[\mu_\alpha X^{(\alpha)}, \sigma_\beta X^{(\beta)}\right] \;&=\; \left(\mu_\alpha \left(\nabla_i \sigma_\beta\right) X^{(\alpha)}(\psi_i)\right) X^{(\beta)} \;+ \\ &-\; \left(\sigma_\beta \left(\nabla_i \mu_\alpha\right) X^{(\beta)}(\psi_i)\right) X^{(\alpha)} \;+\; \left(\mu_a \sigma_\beta\right) \left[X^{(\alpha)}, X^{(\beta)}\right] \;. \end{aligned} \tag{27}$$

Remark. Notice that when $X^{(\beta)} = X_A$, by definition $X^{(\beta)}(\psi_i) = 0$, and

$[X^{(\alpha)}, X^{(\beta)}] = 0$; thus the corresponding subalgebra $\mathcal{X}_\beta$ is always an abelian ideal in $\mathcal{G}$. $\odot$

Remark. Note also that, as obvious from the formula (27) above, the direct sum of subalgebras $\mathcal{X}_{\alpha_1} \oplus ... \oplus \mathcal{X}_{\alpha_s}$ is a subalgebra in $\mathcal{G}$ if and only if $\{X^{(\alpha_1)}, ..., X^{(\alpha_s)}\}$ span a subalgebra in G. $\odot$

Assume now that $\mathcal{G}$ is quasi-linear, and that we are able to determine a sequence of subalgebras $\mathcal{F}_p \subseteq \mathcal{G}$, each of them being the direct sum of $\mathcal{X}_\alpha$ subalgebras, such that $\mathcal{F}_0 = \mathcal{G}$ and

$$[\, \mathcal{G} \,,\, \mathcal{F}_p \,] \;=\; \mathcal{F}_{p+1} \;; \tag{28}$$

if this terminates in zero we say that $\mathcal{G}$ has a *quasi-nilpotent* structure. Notice that the factor algebras $\Gamma_p := \mathcal{F}_p / \mathcal{F}_{p+1}$ are in general *not* abelian.

By the above remark, $\mathcal{G}$ can have a (nontrivial) quasi-nilpotent structure only if G is nilpotent. The chain of subalgebras $\mathcal{F}_p \subset \mathcal{G}$ can then be read off the descending central series G_p of G; recall that the factor algebras $\gamma_p = G_p / G_{p+1}$ for this are abelian. The nonabelian subalgebras Γ_p introduced above are therefore modules over $\mathcal{I}(A)$ generated by abelian subalgebras γ_p of G.

Assume now $\mathcal{G}$ is quasi-nilpotent. In this case we can first work with generators in Γ_1 and simplify terms in Γ_1 (e.g. by following the PRF algorithm within the set Γ_1; this allows to work with more familiar projection and homological equations than if setting the problem in a completely Lie algebraic framework), then consider generators in Γ_2 and simplify the corresponding terms being guaranteed that Γ_1 terms are not changed, and so on.

We will call the reduced normal form obtained in this way the *Lie renormalized form* (LRF); the Lie renormalization is thus a blend of Lie algebraic reduction along the chain of subalgebras identified by (28) and of PRF reduction in each of the Γ_α subsets.

Remark. In this case we are – roughly speaking – just using the nilpotent structure of (the finite dimensional group) G, rather than the one of (the infinite dimensional algebra) $\mathcal{G}$. $\odot$

Remark. Needless to say, this approach is particularly convenient when the Γ_p are generated by a single element of G; it can however be applied also in other situations. $\odot$

Let us briefly discuss why we have considered the quasi-linear case rather than the general one. It should indeed be clear that the key feature is the assumption that we are able to identify $\mathcal{F}_p$ satisfying (28), so that this kind of approach is not confined to the quasi-linear case. However, recall that we have to consider smooth vector fields as generators of our Lie-Poincaré changes of coordinates, and that at each step we have to deal with smooth (partially renormalized) vector fields. As these are only a subset of $\mathcal{X}_1 \oplus ... \oplus \mathcal{X}_d$ (and one identified by analytic, not geometric, conditions), checking these conditions at each step will in general be a prohibitive task. The problem is not present for quasi-linear normal forms, when we deal with the full module $\mathcal{I}_*(A) \otimes G$.

Thus, in practice, the LRF approach is not always applicable; however, when it can be applied the computations required are particularly simple [36].

We also stress that the LRF (like the PRF) for a given system is in general not unique.

We stress that the quasi-nilpotent structure is met in applications: e.g., it applies to any nontrivial two-dimensional case and several three-dimensional ones [36]. More generally, it always (but not only) applies when there is only one basic invariant. We consider now again a simple example; for other explicit ones see [36].

Example. Let us consider again the case considered at the end of the previous section. Proceeding as discussed in this section, we obtain the LRF

$$\dot{w} \;=\; Aw \;+\; (r^{2\mu}\alpha_\mu + r^{4\mu}\alpha_{2\mu})\, w \;+\; \left(\sum_{k=\nu}^{\mu} r^{2k}\beta_k\right) Aw$$

where the sum should be considered as zero for $\mu < \nu$. Notice that now we get a finite dimensional reduced normal form in all cases.

6 Filtering of Lie algebras

I will finally briefly recall (and reformulate, hopefully without distorting it to such an extent to make it completely different from the original version) the general approach employing the Lie algebraic structure of $\mathcal{G}$ (recall this included nonlinear fields only) proposed by Broer and Takens [17, 18], and further developed by other authors.

The Broer procedure is always applicable, and is able produce a unique reduced normal form [6, 7]. Unfortunately, implementing it in practice seems to be quite hard, and this approach has been applied only to systems of very low dimension. On the other side, the ideas put forward by Broer are at the basis of many approaches – including the one depicted in previous sections – to the problem of further reduction of normal forms.

One can consider in general terms the descending central series of $\mathcal{G}$ [49, 54]. We recall this is defined by $\mathcal{G}_0 = \mathcal{G}$ and[14] $\mathcal{G}_{k+1} = [\mathcal{G}, \mathcal{G}_k]$; as well known the factor algebras $\Gamma_k = \mathcal{G}_k/\mathcal{G}_{k+1}$ are abelian. It is then possible to eliminate terms by inner automorphisms of $\mathcal{G}$ (that is, by acting on $\mathcal{G}$ with vector fields in $\mathcal{G}$) proceeding along $\mathcal{G}_k$, i.e. filtering the Lie algebra $\mathcal{G}$. This is relevant to our problem since Lie-Poincaré transformations always reduce to inner operations on $\mathcal{G}$.

This approach is, of course, particularly convenient when $\mathcal{G}$ is nilpotent; actually if $\mathcal{G}_k$ is the descending central series for $\mathcal{G}$, we have $\mathcal{G}_k \cap V_m = \emptyset$ for $m < k$. We can reduce to the more familiar case of algebras nilpotent of finite order N by considering only the projections of $\mathcal{G}_k$ and Γ_k to the linear space $V^N = V_1 \oplus ... \oplus V_N \subset V$.

Let us now briefly describe the implementation of the Broer renormalization procedure in the language employed in the present paper. We consider

[14] Needless to say, $\mathcal{G}_k$ is *not* the set of fields in $\mathcal{G}$ homogeneous of some degree in the x!

the sequence of abelian factor algebras $\Gamma_k = \mathcal{G}_k/\mathcal{G}_{k+1}$, and denote by χ_k the projection from $\mathcal{G}$ to Γ_k.

We will produce a sequence of vector fields $W_*^{(j)}$, each being the result of the first j steps of further normalization. Let η_j be the operator on $\mathcal{G}$ defined by $\eta_j(H) = e^H W_*^{(j)} e^{-H}$; let π_j be the operator of projection from $\mathcal{G}$ to the range of η_j. We can reduce the normal form $W = W_*^{(0)}$ as follows.

As the first step, consider a $H^{(0)} \in \mathcal{G}$ and require that $\widetilde{W} = W_*^{(1)} := \eta_0(H)$ is such that $\pi_0[\chi_0\eta_0(H^{(0)})] = 0$. This determines (non uniquely) $H^{(0)}$, and produces a $W_*^{(1)}$.

At further steps, we have the same setting; the "homological equations" on Lie algebras to be solved at each step will be

$$\pi_j \left[\chi_j \left(\eta_j(H^{(j)}) \right) \right] = 0 ; \tag{29}$$

this determines $H^{(j)}$. Each $W_*^{(j+1)}$ is then determined as

$$W_*^{(j+1)} = \eta_j(H^{(j)}) := \exp[H^{(j)}] \exp[W_*^{(j)}] \exp[-H^{(j)}] . \tag{30}$$

Acknowledgements

The first version of this paper was written while I was a guest of the Theory Group in the Dipartimento di Fisica, Università di Roma; I would like to thank them, and in particular Toni Degasperis, for their continuing and always kind hospitality. The paper has benefitted from remarks by an unknown referee and by the critical readings by G. Cicogna and S. Walcher, whom I warmly thank.

References

[1] A. Algaba, E. Freire and E. Gamero, "Hypernormal forms for equilibria of vector fields. Codimension one linear degeneracies", *Rocky Mount. J. Math.* **29** (1999), 13-45

[2] D. Arnal, M. Ben Ammar and G. Pinczon, "The Poincaré-Dulac theorem for nonlinear representations of nilpotent Lie algebras", *Lett. Math. Phys.* **8** (1984), 467-476

[3] V.I. Arnold, *Geometrical methods in the theory of ordinary differential equations*, Springer, Berlin 1983

[4] V.I. Arnold, *Mathematical Methods of Classical Mechanics*, Springer, Berlin 1983, 1989

[5] V.I. Arnold and Yu.S. Il'yashenko, *Ordinary differential equations*; in: Encyclopaedia of Mathematical Sciences vol. 1 – Dynamical Systems I, (D.V. Anosov and V.I. Arnold eds.), pp. 1-148, Springer, Berlin 1988

[6] A. Baider, "Unique normal forms for vector fields and hamiltonians", *J. Diff. Eqs.* **78** (1989), 33-52

[7] A. Baider and R.C. Churchill, "Uniqueness and non-uniqueness of normal forms for vector fields", *Proc. Royal Soc. Edinburgh* **108A** (1988), 27-33; "Unique normal forms for planar vector fields", *Math. Z.* **199** (1988), 303-310

[8] A. Baider and J. Sanders, "Further reduction of the Takens-Bogdanov normal form", *J. Diff. Eqs.* **99** (1992), 205-244; "Unique normal forms: the nilpotent hamiltonian case", *J. Diff. Eqs.* **92** (1991), 282-304

[9] D. Bambusi and G. Gaeta eds., *Symmetry and Perturbation Theory* (proceedings Torino workshop 1996), C.N.R., Roma 1998

[10] G.R. Belitskii, "Equivalence and normal forms of germs of smooth mappings", *Russ. Math. Surv.* **33** (1978), 107-177

[11] G.R. Belitskii, "Invariant normal forms of formal series", *Funct. Anal. Appl.* **13** (1979), 46-67

[12] G.R. Belitskii, "Normal forms relative to the filtering action of a group", *Trans. Moscow Math. Soc.* **40:2** (1981), 1-39

[13] G.R. Belitskii, "Local normal forms of C^∞ vector fields", paper in this issue

[14] G.R. Belitskii and A.Y. Kopanskii, "Equivariant Sternberg theorem", preprint *mp-arc 00-54*; "Sternberg theorem for equivariant hamiltonian vector fields", *mp-arc 00-449*; "Smooth normalization of reversible vector fields", *mp-arc 99-154*

[15] G. Benettin, L. Galgani and A. Giorgilli, "A proof of the Kolmogorov theorem on invariant tori using canonical transformations defined by the Lie method", *Nuovo Cimento B* **79** (1984), 201-223

[16] H.W. Broer, *Bifurcations of singularities in volume preserving vector fields*, Ph.D. Thesis, Groningen 1979

[17] H.W. Broer, "Formal normal form theorems for vector fields and some consequences for bifurcations in the volume preserving case", in *Dynamical systems and turbulence*, D.A. Rand and L.S. Young eds., *Lect. Notes Math.* **898**, Springer, Berlin 1981

[18] H.W. Broer and F. Takens, "Formally symmetric normal forms and genericity", *Dynamics Reported* **2** (1989), 39-59

[19] A.D. Bruno, *Local methods in the theory of differential equations*, Springer, Berlin 1989; "Analytical form of differential equations I & II", *Trans. Moscow Math. Soc.* **25** (1971), 131-288; and **26** (1971), 199-239

[20] A.D. Bruno, *Power geometry in algebraic and differential equations*, North-Holland, Amsterdam 2000

[21] A.D. Bruno and S. Walcher, "Symmetries and convergence of normalizing transformations", *J. Math. Anal. Appl.* **183** (1994), 571-576

[22] K.T. Chen, "Equivalence and decomposition of vector fields about an elementary critical point", *Am. J. Math.* **85** (1963), 693-722

[23] G. Chen and J. Della Dora, "Further reduction of normal forms for dynamical systems", *J. Diff. Eqs.* **166** (2000), 79-106

[24] L.O. Chua and H. Kokubu, "Normal forms for nonlinear vector fields. Part I: theory", *IEEE Trans. Circ. Syst.* **35** (1988), 863-888 "Normal forms for nonlinear vector fields. Part II: applications", *IEEE Trans. Circ. Syst.* **36** (1989), 51-70

[25] G. Cicogna, "Symmetries of dynamical systems and convergent normal forms", *J. Phys. A* **28** (1995), L179-L182; "On the convergence of the normalizing transformation in the presence of symmetries", *J. Math. Anal. Appl.* **199** (1996), 243-255; "Convergent normal forms of symmetric dynamical systems", *J. Phys. A* **30** (1997), 6021-6028; "Multiple-periodic bifurcation and resonance in dynamical systems", *Nuovo Cimento B* **113** (1998), 1425-1430

[26] G. Cicogna and G. Gaeta, *Symmetry and perturbation theory in nonlinear dynamics*, Springer, Berlin 1999

[27] G. Cicogna and G. Gaeta, "Nonlinear symmetries and normal forms", pp. 3-22 in [29]

[28] G. Cicogna and S. Walcher, "Symmetry and normal forms", paper in this issue

[29] A. Degasperis and G. Gaeta eds., *Symmetry and Perturbation Theory II* (proceedings Roma workshop 1998), World Scientific, Singapore 1999

[30] M. Demazure, *Bifurcation and catastrophes*, Springer, Berlin 2000

[31] A. Deprit, "Canonical transformations depending on a small parameter", *Cel. Mech.* **1** (1969), 12-30

[32] H. Dulac, "Solution d'un systéme d'équations différentielles dans le voisinage des valeurs singuliéres", *Bull. Soc. Math. France* **40** (1912), 324-383

[33] C. Elphick, E. Tirapegui, M. Brachet, P. Coullet and G. Iooss, "A simple global characterization for normal forms of singular vector fields", *Physica D* **29** (1987), 95-127; addendum, *Physica D* **32** (1988), 488

[34] E. Forest and D. Murray, "Freedom in minimal normal forms", *Physica D* **74** (1994), 181-196

[35] G. Gaeta, "Reduction of Poincaré normal forms", *Lett. Math. Phys.* **42** (1997), 103-114; "Poincaré renormalized forms", *Ann. I.H.P. (Phys. Theo.)* **70** (1999), 461-514

[36] G. Gaeta, "Poincaré renormalized forms and regular singular points of vector fields in the plane", preprint *math-ph/0101022*; "Algorithmic reduction of Poincaré-Dulac normal forms and Lie algebraic structure", *Lett. Math. Phys.* **57** (2001), 41-60

[37] G. Gaeta, "Poincaré normal forms and compact simple Lie groups" (revised version), preprint *math-ph/0106016*

[38] A. Giorgilli, "Rigorous results on the power expansions for the integrals of a Hamiltonian system near an elliptic equilibrium point", *Ann. I.H.P.* **48** (1988), 423-439; A. Giorgilli and U. Locatelli, "On classical series expansions for quasi-periodic motions", *Math. Phys. El. J.* **3** (1997); A. Celletti, A. Giorgilli, and U. Locatelli, "Improved estimates on the existence of invariant tori for Hamiltonian systems", *Nonlinearity* **13** (2000), 397-412

[39] A. Giorgilli, "Small denominators and exponential stability: from Poincar to the present time", *Rend. Sem. Mat. Fis. Milano*, to appear; A. Giorgilli and U. Locatelli, "Kolmogorov theorem and classical perturbation theory", *ZAMP* **48** (1997), 220-261

[40] A. Giorgilli, "On the problem of stability for near to integrable Hamiltonian systems", in: Proceedings of the International Congress of Mathematicians Berlin 1998, Vol. III, *Documenta Mathematica, extra volume ICM 1998* (1998), 143-152

[41] P. Glendinning, *Stability, instability and chaos: an introduction to the theory of nonlinear differential equations*, Cambridge University Press, Cambridge 1994

[42] T. Gramchev and M. Yoshino, "Rapidly convergent iterative methods for simultaneous normal forms of commuting maps", *Math. Z.* **231** (1999), 745-770; T. Gramchev, "Simultaneous normal forms of perturbations of vector fields on tori with zero order pseudodifferential operators", pp. 187-195 in [29]; M. Yoshino, "Simultaneous normal forms of commuting maps and vector fields", pp. 287-294 in [29]

[43] J. Guckenheimer and P. Holmes, *Nonlinear oscillations, dynamical systems, and bifurcation of vector fields*, Springer, Berlin 1983

[44] P. Hartman, *Ordinary differential equations*, Wiley, New York 1967

[45] Y. Il'yashenko, "Dulac's memoir 'On limit cycles' and related problems of the local theory of differential equations", *Russ. Math. Surv.* **40:6** (1985), 1-49

[46] Y. Il'yashenko and S. Yakovenko, "Finitely-smooth normal forms of local families of diffeomorphisms and vector fields", *Russ. Math. Surv.* **46** (1991), 1-25

[47] G. Iooss and M. Adelmeyer, "Topics in bifurcation theory and applications, World Scientific, Singapore 1992

[48] H. Ito, "Convergence of the Birkhoff normal form for integrable systems", *Comm. Math. Helv.* **64** (1989), 412-461; "Integrability of hamiltonian systems and Birkhoff normal forms in the simple resonance case", *Math. Ann.* **292** (1992), 411-444

[49] A.A. Kirillov, *Elements of the theory of representations*, Springer 1984

[50] H. Kokubu, H. Oka and D. Wang, "Linear grading functions and further reduction of normal forms", *J. Diff. Eqs.* **132** (1996), 293-318

[51] M. Kummer, "How to avoid secular terms in classical and quantum mechanics", *Nuovo Cimento B*, **1** (1971), 123-148; "On resonant nonlinearly coupled oscillators with two equal frequencies", *Comm. Math. Phys.* **48** (1976), 53-79

[52] L.M. Markhashov, "On the reduction of differential equations to the normal form by an analytic transformation" *J. Appl. Math. Mech.* **38** (1974), 738-740

[53] J-C. van der Meer, *The Hamiltonian Hopf bifurcation*; Lecture Notes in Mathematics **1160**, Springer, Berlin 1985

[54] M.A. Naimark and A.I. Stern, *Theory of group representations*, Springer 1982

[55] V.A. Pliss, "On the reduction of an analytic system of differential equations to linear form", *Diff. Eqs.* **1** (1965), 153-161

[56] H. Rüssmann, "Über das verhalten analytischer hamiltonscher differentialgleichungen in der nahe einer gleichgewichtslosung", *Math. Ann.* **154** (1964), 285-300

[57] K. Siegel and J.K. Moser, *Lectures on Celestial Mechanics*, Springer 1971; reprinted in "Classics in Mathematics", Springer 1995

[58] S. Sternberg, "On the local structure of local homeomorphisms of euclidean n-space", *Amer. J. Math.* **80** (1958), 623-631

[59] L. Stolovitch, "Singular complete integrability", *Publ. Math. I.H.E.S.* **91** (2000), 134-210

[60] L. Stolovitch, "Normalisation holomorphe d'algébres de type Cartan de champs de vecteurs holomorphes singuliers", *C.R. Acad. Sci. (Paris) I* **330** (2000), 121-124

[61] F. Takens, "Singularities of vector fields", *Publ. Math. I.H.E.S.* **43** (1974), 47-100

[62] S. Ushiki, "Normal forms for singularities of vector fields", *Jap. J. Appl. Math.* **1** (1984), 1-34

[63] F. Verhulst, *Nonlinear differential equations and dynamical systems*, Springer, Berlin 1989, 1996

[64] J. Vey, "Sur certains systémes dynamiques séparables", *Am. J. Math.* **100** (1978), 591-614; "Algébres commutatives de champs de vecteurs isochores", *Bull. Soc. Math. France* **107** (1979), 423-432

[65] S. Walcher, "On differential equations in normal form", *Math. Ann.* **291** (1991), 293-314

[66] S. Walcher, "On transformation into normal form", *J. Math. Anal. Appl.* **180** (1993), 617-632

[67] S. Walcher, "Orbital symmetries of first order ODEs", pp. 96-113 in [29]

[68] S. Walcher, "On the Poincaré problem", *J. Diff. Eqs.* **166** (2000), 51-78; "Plane polynomial vector fields with prescribed invariant curves", *Proc. Royal Soc. Edinb.* **130A** (2000), 633-649

[69] S. Walcher, "On convergent normal form transformation in the presence of symmetries", *J. Math. Anal. Appl.* **244** (2000), 17-26